REFERENCE GUIDE

Reference Guide

Aimovig® (AIM-oh-vig) (erenumab-aooe)

Side 1


injection, for subcutaneous use

Single-Dose Prefilled SureClick $^{\scriptsize{(8)}}$ Autoinjector

140 mg/mL

Read all instructions in carton before use.

Questions? Call 1-800-77-AMGEN (1-800-772-6436)



Side 2	Reference Guide Aimovig® (AIM-oh-vig) (erenumab-aooe) injection, for subcutaneous use Single-Dose Prefilled SureClick® Autoinjector 140 mg/mL		Manufactured by: Amgen, Inc. Thousand Oaks, CA 91320 © 2020-2022 Amgen Inc. All rights reserved. This printed material is recyclable. <pre><part number=""> Revised: 8/2022 v3</part></pre>	
Stretch or pinch your injection site to create a firm surface.		Put the yellow safety guard (needle inside) on your skin at 90 degrees.	Firmly push down the autoinjector onto your skin until it stops moving.	Keep pushing down on your skin for about 15 seconds.
3 Stretch metho		4	When you are ready to inject, press the gray start button. You will hear a click.	6 15 seconds
	or	Yellow safety guard (needle inside)	"click"	Window turns from clear to yellow when the injection is done. "click"
Pinch metho	d:			